



Food for Thought ...

Can Case Study Approaches Speed Implementation of the NRC Report: "Toxicity Testing in the 21st Century: A Vision and a Strategy?"

Melvin E. Andersen¹, Harvey J. Clewell, III¹, Paul L. Carmichael², and Kim Boekelheide³

¹The Institute for Chemical Safety Sciences, The Hamner Institutes for Health Sciences, Research Triangle Park, NC, USA; ²Safety and Environmental Assurance Centre, Unilever, Sharnbrook, Bedford, UK;

³Department of Pathology and Laboratory Medicine, Brown University, Providence, RI, USA

Summary

The 2007 report "Toxicity Testing in the 21st Century: A Vision and a Strategy" argued for a change in toxicity testing for environmental agents and discussed federal funding mechanisms that could be used to support this transformation within the USA. The new approach would test for *in vitro* perturbations of toxicity pathways using human cells with high-throughput testing platforms. The NRC report proposed a deliberate timeline, spanning about 20 years, to implement a wholesale replacement of current *in-life* toxicity test approaches focused on apical responses with *in vitro* assays. One approach to accelerating implementation is to focus on well-studied prototype compounds with known toxicity pathway targets. Through a series of carefully executed case studies with four or five pathway prototypes, the various steps required for implementation of an *in vitro* toxicity pathway approach to risk assessment could be developed and refined. In this article, we discuss alternative approaches for implementation and also outline advantages of a case study approach and the manner in which the case studies could be pursued using current methodologies. A case study approach would be complementary to recently proposed efforts to map the human toxome, while representing a significant extension toward more formal risk assessment compared to the profiling and prioritization approaches inherent in programs such as the EPA's ToxCast effort.

Keywords: toxicity testing, case study approach, toxicity pathway

1 Background

The U.S. National Research Council report on "Toxicity Testing in the 21st Century: A Vision and a Strategy" (NRC, 2007; Krewski et al., 2010) argued for a large-scale shift in toxicity testing. Testing protocols for 21st century toxicology should focus on mechanistic *in vitro* assays, typically using human cells in a high-throughput context. The *in vitro* tests would measure perturbations of "toxicity pathways." Perturbations of these key biological signaling pathways are expected to be associated with specific adverse responses when a high level of perturbation persists *in vivo* for sufficient durations. The NRC report has been extensively discussed in the toxicology and risk assessment communities (Andersen and Krewski, 2009; Krewski et al., 2009). The dialog has highlighted four key questions about a shift to *in vitro* tests for assessing risks of chemicals (Andersen and Krewski, 2010): (1) how will we define adversity from *in*

vitro tests; (2) how will the *in vitro* test results be used to predict expected outcomes in animals and people who come in contact with the test compounds; (3) how will regulatory agencies set exposure standards for human populations based on *in vitro* test results; and (4) how will authorities accustomed to the current whole animal testing procedures develop the confidence required to move to another platform for testing and risk assessment. These questions squarely capture the core challenges that need attention in order to develop 21st century toxicology for both toxicity testing and for using these test results for human health risk assessment. The NRC report suggested a process in which incremental advances in key technologies would all have to be completed before making a wholesale transition to *in vitro* test-based risk assessment platforms. Another suggestion is to use case studies with prototype pathways and compounds to accelerate implementation of the Toxicity Testing in the 21st Century (TT21C) paradigm by developing working



examples of the necessary methodologies. Case studies could show each component of this new paradigm in action and effectively probe Questions 1 through 3 noted above. This article begins with considerations of approaches to implementing the NRC's TT21C vision and then considers the process by which case studies might be pursued.

2 Developing the knowledge base for testing

Many of the "Food for thought ..." articles in ALTEX and the dialog regarding reinvention of toxicity testing associated with t⁴ – the transatlantic think tank for toxicology (Rovida and Hartung, 2009) – have provided opportunities to discuss initiatives that promise to reshape testing for many commercial products (Hartung, 2008, 2010b; Hartung and Koeter, 2008; Hartung et al., 2009). Test methods and scientific advances required to move toxicity testing into the 21st century are already in place, such as the application of human stem cells for testing (Leist et al., 2008; Kuegler et al., 2010), computational systems biology for modeling response pathways (Alon, 2007), and evidence-based procedures to guide assay validation (Hoffmann and Hartung, 2006). The challenge is both to utilize these new tools appropriately and to integrate them coherently in 21st century testing strategies.

Implementing these tools into a modern *in vitro* testing paradigm will require the development of new knowledge on toxicity pathways and how they interact and function. Conceptually, this new knowledge base can be accumulated either by using what has been called "an unbiased approach," by using a case study approach, or by a combination of these two approaches (Fig. 1). In the unbiased approach, a great deal of attention would be devoted to designing a convenient *in vitro* test system, based on human cells, that would be expected to include as much of the known biological response landscape as possible. For example, the test system would be designed to include specialized functions of differentiated cell types and the pathways mediating paracrine interactions. This test system would likely include complex bioengineered multicellular test platforms and genetically modified cells with an array of differentiated expression patterns, displaying as complete a representation of known biology as possible. Once this test platform is in place, testing would be expected to gradually reveal the key toxicity pathways and their interactions (Fig. 1). While toxicants with "known" effects would be included in the testing program, in this unbiased approach the inherent responses of the test platforms would be used to define points of departure (POD). The advantages of this unbiased approach include its lack of reliance on past animal-based test results, the open-ended nature of the inquiry, and the continual improvements in reliability and predictivity as more testing takes place. Disadvantages include the importance of the test platform itself as a unique model with unknown universal applicability, and the amount of time required to develop a comprehensive understanding of the toxicity pathways and their interactions prior to their use in human risk assessment.

In the case study approach, existing knowledge of specific toxicity pathways would be used to design appropriate test sys-

tems that would then be exercised with model toxicants before exploring the responses of unknown toxicants (Fig. 1). The advantages of this approach include the focus on well understood pathways and responses that build on our current understanding of cell and molecular biology and the rapid applicability of the test system to a defined set of anticipated responses. The disadvantages include the narrow restriction of the inquiry to known biology, which could lead to missing important unknown responses, and the need to develop different test platforms that are specific to each case study area. In this case study approach, the POD in the *in vitro* test system would be informed by the known *in vivo* responses to the model compounds and would not rely solely on the platform responses. This characteristic would allow alignment of results from new *in vitro* approaches with past whole animal and mode-of-action research.

Ideally, implementation of the new toxicity testing paradigm will be achieved using a combination of unbiased and case study approaches (Fig. 1). By combining approaches, the advantages




Fig. 1: The toxicity pathway knowledge base can be developed using an unbiased approach, a case study approach, or a combined approach

The knowledge needed to develop a functioning *in vitro* test system will be acquired over time (left-to-right). This knowledge can be acquired by having a comprehensive test platform that is challenged by numerous toxicants, and over time the interconnections and important response pathways emerge (unbiased approach). In this "unbiased approach," knowledge expands over time as pathway interactions are understood (reflected by the width of the triangle increasing and the shading darkening from left to right). Alternatively, using a case study approach, a specific pathway can be evaluated in detail. While the density of knowledge increases (the shade darkens with time), the knowledge area covered does not change (the width of the knowledge area remains constant). Combining these approaches may be the best way to take advantage of the strengths of each approach; however, projects to demonstrate the applicability of the new toxicity testing paradigm may be most rapidly developed using a case study approach.

of open-ended inquiry with the focused application of expert knowledge gives an opportunity to develop the most complete toxicity assessment capability. In the near term, however, what is most needed is a demonstration of the feasibility of these new approaches and their ability to be both reliable and predictive. These near-term goals, in our opinions, can most effectively be met with demonstration projects using a case study approach.

One important consideration in implementing the test platforms themselves is the difference between the platforms used to develop the deep biological knowledge base of pathway function and the platforms that will ultimately be used for testing. Because of the absence of the requisite knowledge, the assay platforms for studying specific pathways must be designed to yield integrated data on the interacting molecular systems that control cell function, including transcriptomics, transcriptional factor analysis, phosphoproteomics, metabolomics, etc. Time-dependent changes in cellular function will also need to be assessed to develop a familiarity with the important temporal windows of response. The assay systems for routine testing, however, are likely to be much simpler and to have higher throughput, assessing important integrative nodal responses that adequately assess and reflect the functionality of the cell. The more complex systems will be initially developed to assess assay function; the final applications will simply evaluate high-throughput dose-response. As described by Eric Berlow, an ecologist and network theorist, in a recent TED (Technology Entertainment and Design) Talk (see www.ericberlow.net), “simplicity often lies on the other side of complexity.” Berlow makes a distinction between the complex and the complicated, with complexity arising from lots of interactions and details while complication results from an obscuring of the details. During the development of our understanding of toxicity pathways, the details may at first seem overwhelming and unconnected, but ultimately it will be possible to identify those details that are both important and most representative of the overall responses. The case study approach, then, lays many larger issues aside (Question 4 in the list) in an effort to get started and show how *in vitro* toxicity testing information can be used in specific situations. Success with the case studies should refine the process and the nature of specific pathway assays and help provide context for the more unbiased approaches.

3 Doing risk assessments from *in vitro* test results

The manner in which *in vitro* testing results could be used for risk assessment was outlined in generic fashion in the NRC report (NRC, 2007) and in other thought pieces on new directions in toxicity testing (Krewski et al., 2011). These contributions showed the correspondence of individual components of the proposed toxicity testing methods to the risk assessment paradigm from the 1983 NRC report, *Risk Assessment in the Federal Government: Managing the Process* (NRC, 1983). An essay on the manner in which *in vitro* test results will provide measures of adversity (Boekelheide and Campion, 2010) sparked a broader dialog on designing a more explicit process to develop human exposure guidelines from an *in vitro* concentration deemed to




Fig. 2: A schematic showing steps in a toxicity pathway-based risk assessment

Results from the panel of assays identify the pathway targets and generate a point of departure (POD) for the subsequent risk assessment as an *in vitro* concentration. Computational systems biology pathway (CSBP) modeling of circuitry and dynamics for the assay system indicates the expected shape of the dose response at lower doses, leading to a POD. The POD concentration is then converted to an exposure standard through techniques of reverse dosimetry implemented by pharmacokinetic modeling. This step takes advantage of *in vitro-in vivo* extrapolation (IVIVE).

cause an adverse cellular response (Boekelheide and Andersen, 2010; Bhattacharya et al., 2011). A simple, linear diagram of the likely process (Fig. 2) shows steps involved in obtaining an *in vitro* POD and then moving from the POD to a regulatory standard.

Chemical risk assessment in the future would start by having a suite of “validated” *in vitro* assays and determining the pathway assay or assays that had responses at the lowest test concentration. Some of the challenges for validation of *in vitro* methods were discussed in four papers (Boekelheide and Andersen, 2010; Dietrich, 2010; Hartung, 2010a; Jaworska and Hoffmann, 2010) at a symposium, “21st Century Validation Strategies for 21st Century Tools,” held last July at the Johns Hopkins University. Most of these *in vitro* assays would be pathway specific, with a few that had more broad coverage of cellular responses. The pathway specific assays would have already been validated using positive controls known to affect pathway responses, such as 17- β estradiol for an estrogenic pathway assay or ionizing radiation for a DNA-damage pathway, before use in general testing. The NRC report talked about using high-throughput assays capable of testing hundreds or thousands of compounds very quickly. The goal is not simply to achieve high-throughput but also to have suitable *in vitro* assays that can be run rapidly over a wide range of test concentrations. Testing over a wide range



of concentrations is at the heart of quantitative high-throughput screening (q-HTS). With q-HTS, the empirical dose response curve for the assays would span multiple orders of concentration (Inglese et al., 2006).

From the results of a suite of toxicity pathway tests, the pathways affected at the lowest treatment concentrations would be identified. Computational systems biology pathway (CSBP) models, defined in the process of assay development and validation, would take pathway multi-point dose-response curves and predict transitions from sub-threshold doses to doses causing adaptive changes in the pathway function to doses expected to have adverse consequences. These changes in response patterns with increasing levels of perturbations are referred to as dose-dependent transitions (Slikker et al., 2004a,b). The *in vitro* pathway assay would provide the dose response information for assessing adversity. CSBP models for the assay would inform the low dose extrapolation to give the POD. Then, together with policy considerations, such as sensitive populations, response variability in a diverse population, etc., the *in vitro* adverse concentration would be adjusted to give an acceptable human plasma concentration. The last step in this process is estimating the *in vivo* human exposure expected to produce the *in vitro* concentration. This extrapolation relies on technologies referred to as *in vitro*-*in vivo* extrapolation – IVIVE (Shiran et al., 2006; Gibson and Rostami-Hodjegan, 2007) – and reverse dosimetry (Clewell et al., 2008) – and would have to include considerations of variability in expected response in different populations, co-exposures in mixtures, and other factors.

In vitro to *in vivo* extrapolation (IVIVE) estimates the environmental exposures to a chemical that could produce target tissue exposures in humans that are equivalent to those concentrations that are associated with effects in an *in vitro* toxicity test. Through a combination of quantitative structure property relationship (QSPR) modeling, physiologically based pharmacokinetic (PBPK) modeling, and collection of *in vitro* data on metabolism, transport, binding, etc., IVIVE provides estimates of the likelihood of harmful effects from expected environmental exposures. Generic IVIVE approaches have measured metabolic clearance in primary hepatocytes, plasma protein binding and estimates of renal filtration extrapolated from protein binding. These data were input to a population-based *in vitro*-to-*in vivo* extrapolation program (SIMCYP™) for estimating the human oral equivalent dose necessary to produce a steady-state *in vivo* concentration equivalent to *in vitro* AC₅₀ (concentration at 50% of maximum activity) and LEC (lowest effective concentration) values from the ToxCast data (Rotroff et al., 2010).

4 Developing assays

Human cells in culture

The NRC report favored assays using human cells, cell-lines or organotypic cell cultures in order for the toxicity pathway testing to be more representative of expected effects of test compounds on human biology. In designing pathway assays, there will be compromises about available cells, the nature of pathway assay read-outs, and the presence of pathway components in specific

cells or cell lines. Some assay development could be done in cell lines, but the goal would be to have the final suite of pathway assay testing conducted in human cells. This vision looks more accessible in 2011 than it did in 2007 due to the advances made in using tissue-specific stem cells rather than having to harvest human cells from donors. For instance, hepatoblasts can be differentiated from liver stem cell precursors, maintained, then grown and differentiated into primary hepatocytes for testing in cultures (Wang et al., 2010). As noted in the discussion of unbiased and case study approaches, the development of these assays would require developing a deep biological knowledge of the networks, nodes, and dynamics of pathway responses to test compounds.

Assays then would be developed first with positive control test compounds to illuminate the range of responses associated with different degrees of severity of effect. To illustrate the case study process, we describe approaches to assay design with the DNA-damage, p53 pathway, and then use the PPAR- α pathway to discuss aspects of network inference and CSBP modeling; these are two toxicity pathways currently being developed as case studies. A schematic for the design of assays for specific toxicity pathways (Fig. 3) needs to capture aspects of design for purpose, validation against pathway structure and pathway dynamics, and ability to use the results for mechanistic dose re-




Fig. 3: Developing pathway assays

The steps shown here constitute the process of assuring that pathway assays are fit for purpose, i.e., for identifying activity in a specific pathway, assessing the structure and dynamics of the overall signaling network, and providing computational systems biology pathway models to assist dose-response modeling of assay results and low-dose extrapolations. The process of pathway development is fundamental to their use as part of a testing suite for risk assessment, as shown in Figure 2.

sponse assessment through computational pathway models. Assays used for routine testing would have to first be “validated” through these steps.

DNA-damage assays

Our studies with the DNA-damage related pathways represent an on-going, productive collaboration between Unilever and The Hamner. For this pathway, we are using high content imaging and flow cytometry methods to visualize and quantify markers of DNA-damage (e.g., H2AX and TP53 binding to DNA and micronucleus formation) and activation of the p53 response pathway, including p53, phosphorylated-p53, p21 expression and cell cycle arrest (Lahav et al., 2004; Bryce et al., 2008; van Attikum and Gasser, 2009). Currently, we are using a human osteosarcoma cell line, HT-1080 (Sun et al., 2011). Although this is a cell line rather than a primary human cell, it has functioning native p53 (Benchimol et al., 1982). The goal in this project is to develop a suite of assays at the cellular level that link DNA-damage, pathway responses to DNA-damage, and ultimately mutation. Initially, the responses are modeled using empirical dose response models, including models that test for thresholds (Lutz and Lutz, 2009). These empirical dose-response models differentiate the doses causing specific responses, provide evidence for thresholds, show relative potencies for the multiple responses, and guide CSBP models for DNA-damage that provide a mechanistic basis for these dose-dependent transitions.

Finding the components of the pathways and networks

A recent “Food for thought ...” article (Hartung and McBride, 2011) discussed mapping the human toxome and distinguished among definitions of nodes, pathways, and networks. These case studies are designed to evaluate specific toxicity pathways (in this case specific targets with known biological modes-of-action), but application in principle requires assessing the structure of the signaling network in which the pathway is embedded. Our second case study with the PPAR- α receptor focuses on network inference and computational modeling of the dynamic network. Experimental and bioinformatic tools for inferring structures of biological networks are now widely available (Sachs et al., 2002; di Bernardo et al., 2005; Woolf et al., 2005; Shen et al., 2011), but these tools have not yet been used to examine dose-response or dose-dependent transitions across dose regions within the networks. Dose-response information, both *in vivo* and *in vitro*, is already available that would allow network inference for some pathways, e.g., estrogen receptor(ER) or aryl hydrocarbon receptor (AhR) function (Fertuck et al., 2003; Kwekel et al., 2005; Naciff et al., 2009).

Using GW 7647, a specific PPAR- α agonist, these studies in primary hepatocytes from humans and rats have focused on gene expression and transcription factor binding after treatment (Woods et al., 2011). The study design follows that used to establish a cannabinoid receptor network (Bromberg et al., 2008b). Other inference methods (Shen et al., 2011) apply Bayesian approaches and combine gene expression, ChIP-chip or ChIP-seq methods for transcription factor binding and DNA-sequence analysis for locating DNA-response elements throughout the genome. The expected output for each pathway would be a




Fig. 4: A schematic for a “developmental network” that controls receptor-mediated signaling

The network is controlled by several incoherent (iFFL) and coherent feed forward loops (cFFL) with nodes to either activate (arrows) or repress (blunt lines) gene expression through critical signaling nodes – the Y and Z factors. The goal in network inference is to understand the circuitry of various toxicity pathways at this level of detail in order to describe exposures that are without appreciable effects, doses with activation of early portions of the pathway, and full activation. These dose dependencies are expected to coincide with areas of sub-threshold, adaptive, and adverse perturbations as outlined in the NRC report (NRC, 2007). The proteins designated Z1, Z2, and Z3 would have non-monotonic dose response curves; Y1, Y2, Y3, and Y4 would have monotonically increasing, but sequentially delayed, responses. Based on similar networks previously described (Alon, 2007); the code can be obtained by contacting MEA (MAndersen@thehamner.org).

grouping of nodes connected in a logical pattern leading to a sequential dose-dependent network that accounts for the consequences of pathway activation. The resulting circuit structure is likely to contain sequential nodes controlling gene expression, similar in structure to what has been called a “developmental network” (Alon, 2007). The schematic of a developmental network (Fig. 4) would contain multiple nodes and feed-forward loops, thereby creating a dynamic cascade of sequential activation of sub-components of the network. The cascade behavior of these networks (Landers and Spelsberg, 1992) is expected in turn to show both time and dose dependencies in network function. In addition, high-throughput phosphoproteomic data now provide opportunities to construct extended pathways that



link receptor activation with cellular phenotypes (Samaga et al., 2009; Melas et al., 2011); more limited evaluation of contributions of mitogen-activated protein kinases (MAPKs) to specific phenotypes after pathway activation have shown the differential control of key signaling nodes within overall networks (Bromberg et al., 2008a; Zorina et al., 2010). Integrated systems biology data – gene expression, phosphoprotein alterations, transcription factor analysis – will be needed to provide the dynamic and functional characterization of these assays and their associated toxicity pathways.

CSBP models

For each toxicity pathway, the goal is to have a CSBP model that accounts for the variable response dynamics, depending on the level of activation of the network. A variety of modeling tools are now available and have become widely used in the biomedical engineering community (Aldridge et al., 2006). Our primary efforts with CSBP models are currently focused on p53-mediated DNA-damage and repair networks. High-dose responses of p53-mediated DNA-damage pathways have been examined through iterative experimentation and pathway modeling by the biomedical community for the past 10 years (Lahav et al., 2004; Batchelor et al., 2008, 2009; Loewer et al., 2010). In application of the assays on a routine basis for testing (Fig. 2), network inference and CSBP models would not be determined for each compound studied. These tools are part of the process of assay development, determination of fitness of the assays for purpose, and use a deep understanding of network biology to consider dose-dependent transitions in network activation.

5 Running case studies

A case study approach takes well-studied compounds that are known to affect specific pathways and runs them through a process to assess dose-response and convert the POD to a human exposure standard (Fig. 2). Candidates for these case studies are compounds that have been studied with conventional *in vivo* testing methods and whose mode-of-action has been well-characterized from both the *in vivo* test results and secondary mechanistic research. The mode-of-action framework activities over the past decade could serve as the basis for selecting one or more candidates (Sonich-Mullin et al., 2001; Boobis et al., 2006, 2008; Julien et al., 2009). Case study approaches with prototype compounds requires the availability of appropriate assays for specific cellular toxicity pathways and genomic/bioinformatic tools to infer network structure and to create computational systems biology pathway (CSBP) models. For specific test compounds, IVIVE methods require either chemical or chemical class specific approaches for kinetic modeling to describe expected pharmacokinetics and to support reverse dosimetry. Some initial work has focused on both receptor-mediated toxicity pathways (PAPR- α , ER, AhR, etc.) and pathways associated with chemical reactivity, such as DNA-damage, oxidative stress, and metal stress pathways (Simmons et al., 2009). Receptor-mediated pathways, especially estrogen, androgen, and thyroid hormone networks, are

of great interest because of contemporary focus on endocrine-active compounds, aka endocrine disruptors, and the regulatory requirement for testing these compounds in a tiered approach from limited *in vitro* assays to more complex multi-generational *in vivo* studies. To gain confidence in their applicability, the case studies will require optimization and characterization of the assay systems (Fig. 3), followed by exercising of the overall process with model chemicals and other pathway active compounds (Fig. 2).

6 Conclusions

At least three different approaches to implementation of TT21C have now been articulated: a toxome approach (Hartung and McBride, 2011), the US EPA ToxCast program (<http://www.epa.gov/nct/toxcast/>) together with the associated multi-agency Tox21 initiative (<http://www.epa.gov/nct/Tox21/>), and the case study approach described here. Each of these approaches has strengths and contributes to the overall goal of modernizing toxicity testing. The toxome project would map the entirety of pathways of toxicity in humans; ToxCast/Tox21 seeks to develop ways to predict potential toxicity and to develop a cost-effective approach for prioritizing the thousands of chemicals that need toxicity testing (Collins et al., 2008). The case studies are intended to show the application of a new understanding of toxicity pathways directly for human health risk assessment and to do it quickly. Progress in developing case studies will require careful selection of prototype compounds and prototype pathways. In addition, contributions are necessary across disciplines – assay design, genomics/bioinformatics, computational modeling, pharmacokinetics and human health risk assessment. The case study approach offers some advantages. First, it focuses on developing the generic tools and the processes by which *in vitro* toxicity information will be used for setting regulatory standards in specific instances. Second, it proposes learning by doing. Many key issues relevant to the use of this new information will become apparent by moving ahead with examples rather than worrying over how to make wholesale changes. Two of the authors of this perspective (MEA and HJC, III) can look to the advances in PBPK modeling over the past 30 years. It is no exaggeration to say that the majority of challenges required to implement PBPK modeling to a diverse set of compounds were clearly defined with results from the first two compounds – styrene and methylene chloride (Ramsey and Andersen, 1984; Andersen et al., 1987). History is likely to repeat itself with TT21C. After completing the first two or three pathway case studies, most of the issues will become clear and expansion of the testing to other pathways will be greatly accelerated. We are also very much aware from our experiences that in the process of developing case studies, the schematic of the application of toxicity pathway information for risk assessment (Fig. 2) will likely change. In addition, we expect that development of the specific case studies will also help illuminate the steps that will be needed to go forward with the unbiased approaches (Fig. 1). The greatest benefit, though, is likely to be the ability to accelerate change compared to the



other approaches. Case studies should serve to markedly speed the evolution of toxicology with a corresponding reduction in animal testing (Hartung and Leist, 2008).

References

- Aldridge, B. B., Burke, J. M., Lauffenburger, D. A., and Sorger, P. K. (2006). Physicochemical modelling of cell signalling pathways. *Nat. Cell Biol.* 8, 1195-1203.
- Alon, U. (2007). Network motifs: theory and experimental approaches. *Nat. Rev. Genet.* 8, 450-461.
- Andersen, M. E., Clewell, H. J., 3rd, Gargas, M. L., et al. (1987). A physiologically based pharmacokinetics and the risk assessment process for methylene chloride. *Toxicol. Appl. Pharmacol.* 87, 185-205.
- Andersen, M. E., and Krewski, D. (2009). Toxicity testing in the 21st century: bringing the vision to life. *Toxicol. Sci.* 107, 324-330.
- Andersen, M. E., and Krewski, D. (2010). The vision of toxicity testing in the 21st century: Moving from discussion to action. *Toxicol. Sci.* 117, 17-24.
- Batchelor, E., Mock, C. S., Bhan, I., et al. (2008). Recurrent initiation: A mechanism for triggering p53 pulses in response to DNA damage. *Mol. Cell* 30, 277-289.
- Batchelor, E., Loewer, A., and Lahav, G. (2009). The ups and downs of p53: understanding protein dynamics in single cells. *Nat. Rev. Cancer* 9, 371-377.
- Benchimol, S., Pim, D., and Crawford, L. (1982). Radioimmunoassay of the cellular protein p53 in mouse and human cell lines. *Embo J.* 1, 1055-1062.
- Bhattacharya, S., Zhang, Q., Carmichael, P. L., et al. (2011). Toxicity testing in the 21st century: defining new risk assessment approaches based on perturbation of intracellular toxicity pathways. *PLoS One* 6, e20887.
- Boekelheide, K., and Andersen, M. E. (2010). A mechanistic redefinition of adverse effects – a key step in the toxicity testing paradigm shift. *ALTEX* 27, 243-252.
- Boekelheide, K., and Campion, S. N. (2010). Toxicity testing in the 21st century: using the new toxicity testing paradigm to create a taxonomy of adverse effects. *Toxicol. Sci.* 114, 20-24.
- Boobis, A. R., Cohen, S. M., Dellarco, V., et al. (2006). IPCS framework for analyzing the relevance of a cancer mode of action for humans. *Crit. Rev. Toxicol.* 36, 781-792.
- Boobis, A. R., Doe, J. E., Heinrich-Hirsch, B., et al. (2008). IPCS framework for analyzing the relevance of a noncancer mode of action for humans. *Crit. Rev. Toxicol.* 38, 87-96.
- Bromberg, K. D., Iyengar, R., and He, J. C. (2008a). Regulation of neurite outgrowth by G(i/o) signaling pathways. *Front. Biosci.* 13, 4544-4557.
- Bromberg, K. D., Ma'ayan, A., Neves, S. R., and Iyengar, R. (2008b). Design logic of a cannabinoid receptor signaling network that triggers neurite outgrowth. *Science* 320, 903-909.
- Bryce, S. M., Avlasevich, S. L., Bemis, J. C., et al. (2008). Interlaboratory evaluation of a flow cytometric, high content in vitro micronucleus assay. *Mutat. Res.* 650, 181-195.
- Clewell, H. J., Tan, Y. M., Campbell, J. L., and Andersen, M. E. (2008). Quantitative interpretation of human biomonitoring data. *Toxicol. Appl. Pharmacol.* 231, 122-133.
- Collins, F. S., Gray, G. M., and Bucher, J. R. (2008). Toxicology. Transforming environmental health protection. *Science* 319, 906-907.
- di Bernardo, D., Thompson, M. J., Gardner, T. S., et al. (2005). Chemogenomic profiling on a genome-wide scale using reverse-engineered gene networks. *Nat. Biotechnol.* 23, 377-383.
- Dietrich, D. R. (2010). Courage for simplification and imperfection in the 21st century assessment of “Endocrine disruption”. *ALTEX* 27, 264-278.
- Fertuck, K. C., Eckel, J. E., Gennings, C., and Zacharewski, T. R. (2003). Identification of temporal patterns of gene expression in the uteri of immature, ovariectomized mice following exposure to ethynylestradiol. *Physiol. Genomics* 15, 127-141.
- Gibson, G. G., and Rostami-Hodjegan, A. (2007). Modelling and simulation in prediction of human xenobiotic absorption, distribution, metabolism and excretion (ADME): in vitro-in vivo extrapolations (IVIVE). *Xenobiotica* 37, 1013-1014.
- Hartung, T. (2008). Food for thought ... on alternative methods for cosmetics safety testing. *ALTEX* 25, 147-162.
- Hartung, T., and Koëter, H. (2008). Food for thought ... on food safety testing. *ALTEX* 25, 259-264.
- Hartung, T., and Leist, M. (2008). Food for thought ... on the evolution of toxicology and the phasing out of animal testing. *ALTEX* 25, 91-102.
- Hartung, T., Blaauboer, B., and Leist, M. (2009). Food for thought ... on education in alternative methods in toxicology. *ALTEX* 26, 255-263.
- Hartung, T. (2010a). Evidence-based toxicology – the toolbox of validation for the 21st century? *ALTEX* 27, 253-263.
- Hartung, T. (2010b). Food for thought ... on alternative methods for nanoparticle safety testing. *ALTEX* 27, 87-95.
- Hartung, T., and McBride, M. (2011). Food for thought ... on mapping the human toxome. *ALTEX* 28, 83-93.
- Hoffmann, S., and Hartung, T. (2006). Toward an evidence-based toxicology. *Hum. Exp. Toxicol.* 25, 497-513.
- Inglese, J., Auld, D. S., Jadhav, A., et al. (2006). Quantitative high-throughput screening: a titration-based approach that efficiently identifies biological activities in large chemical libraries. *Proc. Natl. Acad. Sci. USA* 103, 11473-11478.
- Jaworska, J., and Hoffmann, S. (2010). Integrated Testing Strategy (ITS) – Opportunities to better use existing data and guide future testing in toxicology. *ALTEX* 27, 231-242.
- Julien, E., Boobis, A. R., and Olin, S. S. (2009). The key events dose-response framework: a cross-disciplinary mode-of-action based approach to examining dose-response and thresholds. *Crit. Rev. Food Sci. Nutr.* 49, 682-689.
- Krewski, D., Andersen, M. E., Mantus, E., and Zeise, L. (2009). Toxicity testing in the 21st century: Implications for human health risk assessment. *Risk Anal.* 29, 474-479.
- Krewski, D., Acosta, D., Andersen, M., et al. (2010). Toxicity testing in the 21st century: A vision and a strategy. *J. Toxicol. Environ. Health B Crit. Rev.* 13, 51-138.
- Krewski, D., Westphal, M., Al-Zoughool, M., et al. (2011). New directions in toxicity testing. *Annu. Rev. Public Health* 32, 161-178.
- Kuegler, P. B., Zimmer, B., Waldmann, T., et al. (2010). Markers of murine embryonic and neural stem cells, neurons and astro-



- cytes: reference points for developmental neurotoxicity testing. *ALTEX* 27, 17-42.
- Kwekel, J. C., Burgoon, L. D., Burt, J. W., et al. (2005). A cross-species analysis of the rodent uterotrophic program: elucidation of conserved responses and targets of estrogen signaling. *Physiol. Genomics* 23, 327-342.
- Lahav, G., Rosenfeld, N., Sigal, A., et al. (2004). Dynamics of the p53-Mdm2 feedback loop in individual cells. *Nat. Genet.* 36, 147-150.
- Landers, J. P., and Spelsberg, T. C. (1992). New concepts in steroid hormone action: transcription factors, proto-oncogenes, and the cascade model for steroid regulation of gene expression. *Crit. Rev. Eukaryot. Gene Expr.* 2, 19-63.
- Leist, M., Bremer, S., Brundin, P., et al. (2008). The biological and ethical basis of the use of human embryonic stem cells for in vitro test systems or cell therapy. *ALTEX* 25, 163-190.
- Loewer, A., Batchelor, E., Gaglia, G., and Lahav, G. (2010). Basal dynamics of p53 reveal transcriptionally attenuated pulses in cycling cells. *Cell* 142, 89-100.
- Lutz, W. K., and Lutz, R. W. (2009). Statistical model to estimate a threshold dose and its confidence limits for the analysis of sub-linear dose-response relationships, exemplified for mutagenicity data. *Mutat. Res.* 678, 118-122.
- Melas, I. N., Mitsos, A., Messinis, D. E., et al. (2011). Combined logical and data-driven models for linking signalling pathways to cellular response. *BMC Syst. Biol.* 5, 107.
- Naciff, J. M., Khambatta, Z. S., Thomason, R. G., et al. (2009). The genomic response of a human uterine endometrial adenocarcinoma cell line to 17alpha-ethynyl estradiol. *Toxicol. Sci.* 107, 40-55.
- NRC (1983). *Risk assessment in the federal government: managing the process*. Washington, DC: The National Academy Press.
- NRC (2007). *Toxicity testing in the 21st century: A vision and a strategy*. Washington, DC: The National Academies Press.
- Ramsey, J. C., and Andersen, M. E. (1984). A physiologically based description of the inhalation pharmacokinetics of styrene in rats and humans. *Toxicol. Appl. Pharmacol.* 73, 159-175.
- Rotroff, D. M., Beam, A. L., Dix, D. J., et al. (2010). Xenobiotic-metabolizing enzyme and transporter gene expression in primary cultures of human hepatocytes modulated by ToxCast chemicals. *J. Toxicol. Environ. Health B Crit. Rev.* 13, 329-346.
- Rovida, C., and Hartung, T. (2009). Re-evaluation of animal numbers and costs for in vivo tests to accomplish REACH legislation requirements for chemicals – a report by the transatlantic think tank for toxicology (t⁴). *ALTEX* 26, 187-208.
- Sachs, K., Gifford, D., Jaakkola, T., et al. (2002). Bayesian network approach to cell signaling pathway modeling. *Sci. STKE* 2002, pe38.
- Samaga, R., Saez-Rodriguez, J., Alexopoulos, L. G., et al. (2009). The logic of EGFR/ErbB signaling: theoretical properties and analysis of high-throughput data. *PLoS Comput. Biol.* 5, e1000438.
- Shen, C., Huang, Y., Liu, Y., et al. (2011). A modulated empirical Bayes model for identifying topological and temporal estrogen receptor alpha regulatory networks in breast cancer. *BMC Syst. Biol.* 5, 67.
- Shiran, M. R., Proctor, N. J., Howgate, E. M., et al. (2006). Prediction of metabolic drug clearance in humans: in vitro-in vivo extrapolation vs allometric scaling. *Xenobiotica* 36, 567-580.
- Simmons, S. O., Fan, C. Y., and Ramabhadran, R. (2009). Cellular stress response pathway system as a sentinel ensemble in toxicological screening. *Toxicol. Sci.* 111, 202-225.
- Slikker, W., Jr, Andersen, M. E., Bogdanffy, M. S., et al. (2004a). Dose-dependent transitions in mechanisms of toxicity: case studies. *Toxicol. Appl. Pharmacol.* 201, 226-294.
- Slikker, W., Jr, Andersen, M. E., Bogdanffy, M. S., et al. (2004b). Dose-dependent transitions in mechanisms of toxicity. *Toxicol. Appl. Pharmacol.* 201, 203-225.
- Sonich-Mullin, C., Fielder, R., Wiltse, J., et al. (2001). IPCS conceptual framework for evaluating a mode of action for chemical carcinogenesis. *Regul. Toxicol. Pharmacol.* 34, 146-152.
- Sun, B., Daniels, K., Ross, S., et al. (2011). Dose-response evaluation of p53 and H2AX DNA damage response by methyl methanesulfonate, etoposide, and quercetin. *The Toxicologist*, 24.
- van Attikum, H., and Gasser, S. M. (2009). Crosstalk between histone modifications during the DNA damage response. *Trends Cell Biol.* 19, 207-217.
- Wang, Y., Yao, H. L., Cui, C. B., et al. (2010). Paracrine signals from mesenchymal cell populations govern the expansion and differentiation of human hepatic stem cells to adult liver fates. *Hepatology* 52, 1443-1454.
- Woods, C. G., Yarborough, K. M., Pluta, L. J., et al. (2011). Constructing a PPAR α -mediated transcriptional network in primary human and rat hepatocytes. *The Toxicologist*, 78.
- Woolf, P. J., Prudhomme, W., Daheron, L., et al. (2005). Bayesian analysis of signaling networks governing embryonic stem cell fate decisions. *Bioinformatics* 21, 741-753.
- Zorina, Y., Iyengar, R., and Bromberg, K. D. (2010). Cannabinoid 1 receptor and interleukin-6 receptor together induce integration of protein kinase and transcription factor signaling to trigger neurite outgrowth. *J. Biol. Chem.* 285, 1358-1370.

Acknowledgments

MEA and HJC thank the ACC-LRI (Long Range Research Initiative of the American Chemistry Council), Dow Chemical, Unilever, the Exxon Mobil Foundation, and Dow Corning for support of method development and data generation for pursuing case study approaches to bringing the TT21C vision to life more quickly. Support for KB was provided in part by the Superfund Research Program of the National Institute of Environmental Health Sciences at the National Institutes of Health (P42 ES013660).

Correspondence to

Melvin Andersen, PhD
The Institute for Chemical Safety Sciences
The Hamner Institutes for Health Sciences
Six Davis Drive, PO Box 12137
Research Triangle Park, NC 27709-2137, USA
Phone: 919-558-1205 Fax: 919-558-1300
e-mail: mandersen@thehamner.org